


I

(1)

$$f(x) = \begin{cases} \frac{1}{2}x + \frac{1}{2} & (x \leq 1) \\ 2x - 1 & (x > 1) \end{cases}$$



(1)

 $x \leq 1$ のとき

$$\begin{aligned} f(x) - x &= \frac{1}{2}x + \frac{1}{2} - x \\ &= \frac{1}{2} - \frac{1}{2}x \\ &= \frac{1}{2}(1 - x) \leq 0 \end{aligned}$$

 $x > 1$ のとき

$$\begin{aligned} f(x) - x &= 2x - 1 - x \\ &= x - 1 > 0 \end{aligned}$$

よって成立。

(2)

$$a_1 = a \leq 1$$

 $a_n \leq 1$ とすると

$$a_{n+1} = f(a_n) = \frac{1}{2}a_n + \frac{1}{2} \text{ より}$$

$$1 - a_{n+1} = 1 - \left(\frac{1}{2}a_n + \frac{1}{2} \right) = \frac{1}{2}(1 - a_n) \geq 0$$

よって 数学的帰納法により すべての自然数 n に対して $a_n \leq 1$

(3)

(2) より、すべての自然数 n に対して $a_n \leq 1$ なので

$$a_{n+1} = \frac{1}{2}a_n + \frac{1}{2}, \quad a_1 = 1 (\leq 1)$$


$$\Leftrightarrow a_{n+1} - 1 = \frac{1}{2}(a_n - 1)$$

$$a_n - 1 = (a - 1) \left(\frac{1}{2} \right)^{n-1}$$

$$\therefore a_n = 1 - (1 - a) \left(\frac{1}{2} \right)^{n-1}$$

II


(1)

(i) 判別式 : $D > 0$

$$\Leftrightarrow a^2 - 4b > 0 \Leftrightarrow b < \frac{1}{4}a^2$$

(ii) 軸 : $x = -\frac{a}{2} > 0$

$$\Leftrightarrow a < 0$$

(iii) $f(0) = b > 0$ (2) $f(x) = 0$ の解が実数のとき(i) 判別式 : $D \geq 0$

$$\Leftrightarrow b \leq \frac{1}{4}a^2$$

(ii) 軸 : $x = -\frac{a}{2} < 0$


$$\Leftrightarrow a > 0$$

(iii) $f(0) = b > 0$


解が虚数のとき

$$(i) D < 0 \Leftrightarrow b > \frac{1}{4}a^2$$

$$(ii) \text{ 解の実部} : -\frac{a}{2} < 0 \Leftrightarrow a > 0$$

以上より $a > 0, b > 0$ 

(3)


 $f(x) = 0$ の解が実数のとき

解が虚数のとき

(i) 判別式 : $D < 0 \Leftrightarrow b > \frac{1}{4}a^2$

(ii) 解の実部 : $-1 < -\frac{a}{2} < 0$
 $\Leftrightarrow 0 < a < 2$

以上より



III

(1)

偶数の n 枚から 2 枚または素数の n 枚から 2 枚を選べばよい

$$\frac{{}_nC_2 + {}_nC_2}{2{}_nC_2} = \frac{2n(n-1)}{2n(2n-1)} = \frac{n-1}{2n-1}$$

(2)

3 枚のカードの和が偶数となるのは


- (i) 3 枚とも偶数 : ${}_nC_3$ 通り
- (ii) 1 枚が偶数, 2 枚が奇数 : ${}_nC_1 \times {}_nC_2$

の場合なので

$$\begin{aligned} \frac{{}_nC_3 + {}_nC_1 \times {}_nC_2}{2{}_nC_3} &= \frac{\frac{1}{6}n(n-1)(n-2) + n \times \frac{n(n-1)}{2}}{\frac{1}{6}2n(2n-1)(2n-2)} \\ &= \frac{n(n-1)(n-2) + 3n^2(n-1)}{4n(2n-1)(n-1)} \\ &= \frac{n-2+3n}{4(2n-1)} = \frac{2(2n-1)}{4(2n-1)} = \frac{1}{2} \end{aligned}$$

(3)

カードの取り出し方は



$$(1 + 2 + 3 + \cdots + 2n) - n = \frac{2n(2n+1)}{2} - n = 2n^2 \text{ 通り}$$


全体は

$$(2n)^2 - 2n = 2n(2n-1) \text{ 通り} \text{ なので}$$

$$\frac{2n^2}{2n(2n-1)} = \frac{n}{2n-1}$$



IV



(1)

$$\begin{aligned} |\overrightarrow{AB}|^2 &= |\overrightarrow{OA}|^2 + |\overrightarrow{OB}|^2 - 2\overrightarrow{OA} \cdot \overrightarrow{OB} \\ &= 13 + 25 - 2 = 36 \end{aligned}$$

$$\therefore |\overrightarrow{AB}| = 6$$

(2)

$$\begin{aligned} \overrightarrow{OH} &= \overrightarrow{OA} + s\overrightarrow{AB} + t\overrightarrow{AC} \\ &= (1-s-t)\overrightarrow{OA} + s\overrightarrow{OB} + t\overrightarrow{OC} \end{aligned}$$

 $r = 1 - s - t$ とおくと

$$\overrightarrow{OH} = r\overrightarrow{OA} + s\overrightarrow{OB} + t\overrightarrow{OC}, \quad r + s + t = 1$$

OH は△ABC に垂直なので

$$\overrightarrow{OH} \cdot \overrightarrow{AB} = 0 \text{ かつ } \overrightarrow{OH} \cdot \overrightarrow{AC} = 0$$

$$\begin{aligned} &\Leftrightarrow \overrightarrow{OH} \cdot \overrightarrow{OA} = \overrightarrow{OH} \cdot \overrightarrow{OB} = \overrightarrow{OH} \cdot \overrightarrow{OC} \\ &\Leftrightarrow r|\overrightarrow{OA}|^2 + s\overrightarrow{OA} \cdot \overrightarrow{OB} + t\overrightarrow{OA} \cdot \overrightarrow{OC} \\ &\quad = r\overrightarrow{OA} \cdot \overrightarrow{OB} + s|\overrightarrow{OB}|^2 + t\overrightarrow{OB} \cdot \overrightarrow{OC} \\ &\quad = r\overrightarrow{OA} \cdot \overrightarrow{OC} + s\overrightarrow{OB} \cdot \overrightarrow{OC} + t|\overrightarrow{OC}|^2 \\ &\Leftrightarrow 13r + s + t = r + 25s - 11t = r - 11s + 25t \end{aligned}$$

これを解いて

$$\begin{array}{ll} 12r + 12t = 24s & 36s = 36t \\ \Leftrightarrow r + t = 2s & \Leftrightarrow s = t = \frac{1}{3} \\ \Leftrightarrow 1 - s = 2s & \\ \Leftrightarrow s = \frac{1}{3} & \Leftrightarrow r = \frac{1}{3} \end{array}$$

$$\therefore s = t = \frac{1}{3}$$

(3)

$$\begin{aligned}
 |\overrightarrow{OH}|^2 &= \left| \frac{1}{3} (\overrightarrow{OA} + \overrightarrow{OB} + \overrightarrow{OC}) \right|^2 \\
 &= \frac{1}{9} \left\{ |\overrightarrow{OA}|^2 + |\overrightarrow{OB}|^2 + |\overrightarrow{OC}|^2 + 2(\overrightarrow{OA} \cdot \overrightarrow{OB} + \overrightarrow{OB} \cdot \overrightarrow{OC} + \overrightarrow{OC} \cdot \overrightarrow{OA}) \right\} \\
 &= \frac{1}{9} \{13 + 25 + 25 + 2(1 + 1 - 11)\} \\
 &= \frac{1}{9}(63 - 18) = 5
 \end{aligned}$$

$$|\overrightarrow{OH}| = \sqrt{5}$$

$$\begin{aligned}
 |\overrightarrow{AC}|^2 &= |\overrightarrow{OA}|^2 + |\overrightarrow{OC}|^2 - 2\overrightarrow{OA} \cdot \overrightarrow{OC} = 13 + 25 - 2 = 36 \\
 \therefore |\overrightarrow{AC}| &= 6
 \end{aligned}$$

$$\begin{aligned}
 \overrightarrow{AB} \cdot \overrightarrow{AC} &= |\overrightarrow{OA}|^2 - \overrightarrow{OA} \cdot (\overrightarrow{OB} + \overrightarrow{OC}) + \overrightarrow{OB} \cdot \overrightarrow{OC} \\
 &= 13 - 1 - 1 - 11 = 0
 \end{aligned}$$

$$\therefore \triangle ABC = 6 \times 6 \times \frac{1}{2} = 18$$

$$OABC = \frac{1}{3} \times 18 \times 5 = 30$$

V

$$x = \sin t$$

$$y = \cos\left(t - \frac{\pi}{6}\right) \sin t \quad (0 \leq t \leq \pi)$$

(1)

$$\frac{dx}{dt} = \cos t = 0 \quad \therefore t = \frac{\pi}{2}$$

$$\begin{aligned} \frac{dy}{dt} &= -\sin\left(t - \frac{\pi}{6}\right) \sin t + \cos\left(t - \frac{\pi}{6}\right) \cos t \\ &= \cos\left(2t - \frac{\pi}{6}\right) = 0 \end{aligned}$$


$$\therefore 2t - \frac{\pi}{6} = \frac{\pi}{2}, \frac{3}{2}\pi$$

$$2t = \frac{2}{3}\pi, \frac{5}{3}\pi$$

$$t = \frac{\pi}{3}, \frac{5}{6}\pi$$

(2)

t	0		$\frac{\pi}{3}$		$\frac{\pi}{2}$		$\frac{5}{6}\pi$		π
$\frac{dx}{dt}$		+		+	0	-		-	
$\frac{dy}{dt}$		+	0	-		-	0	+	
x	0	\rightarrow	$\frac{\sqrt{3}}{2}$	\rightarrow	1	\leftarrow	$\frac{1}{2}$	\leftarrow	0
y	0	\uparrow	$\frac{3}{4}$	\downarrow	$\frac{1}{2}$	\downarrow	$-\frac{1}{4}$	\uparrow	0
		\nearrow		\searrow		\swarrow		\nwarrow	



(3)

$$y = 0 \text{ となるのは } t = 0, \pi, \frac{2}{3}\pi \quad t = \frac{2}{3}\pi \text{ のとき } x = \frac{\sqrt{3}}{2}$$

よって求める面積は C の $y \leq 0$ の部分を y_- とすると

$$\begin{aligned} - \int_0^{\frac{\sqrt{3}}{2}\pi} y_- dx &= - \int_{\pi}^{\frac{2}{3}\pi} \cos\left(t - \frac{\pi}{6}\right) \sin t \cos t dt \\ &= \int_{\frac{2}{3}\pi}^{\pi} \frac{1}{2} \cos\left(t - \frac{\pi}{6}\right) \sin 2t dt \\ &= \frac{1}{2} \int_{\frac{2}{3}\pi}^{\pi} \frac{1}{2} \left\{ \sin\left(3t - \frac{\pi}{6}\right) + \sin\left(t + \frac{\pi}{6}\right) \right\} dt \\ &= \frac{1}{4} \left[-\frac{1}{3} \cos\left(3t - \frac{\pi}{6}\right) - \cos\left(t + \frac{\pi}{6}\right) \right]_{\frac{2}{3}\pi}^{\pi} \\ &= -\frac{1}{4} \left[\frac{1}{3} \cos\left(3t - \frac{\pi}{6}\right) + \cos\left(t + \frac{\pi}{6}\right) \right]_{\frac{2}{3}\pi}^{\pi} \\ &= -\frac{1}{4} \left\{ \frac{1}{3} \cos\left(3\pi - \frac{\pi}{6}\right) + \cos\left(\pi + \frac{\pi}{6}\right) - \frac{1}{3} \cos\left(2\pi - \frac{\pi}{6}\right) - \cos\left(\frac{2}{3}\pi + \frac{\pi}{6}\right) \right\} \\ &= -\frac{1}{4} \left\{ \frac{1}{3} \cos \frac{5}{6}\pi + \cos \frac{7}{6}\pi - \frac{1}{3} \cos \frac{\pi}{6} - \cos \frac{5}{6}\pi \right\} \\ &= -\frac{1}{4} \left(-\frac{2}{3} \left(-\frac{\sqrt{3}}{2} \right) - \frac{\sqrt{3}}{2} - \frac{1}{3} \cdot \frac{\sqrt{3}}{2} \right) \\ &= -\frac{1}{4} \cdot \frac{\sqrt{3}}{2} \left(\frac{2}{3} - 1 - \frac{1}{3} \right) = \frac{\sqrt{3}}{12} \end{aligned}$$